

Characterization of a Novel Picornavirus Isolate from a Diseased European Eel (*Anguilla anguilla*)

Dieter Fichtner, Anja Philipps, Marco Groth, Heike Schmidt-Posthaus, Harald Granzow, Malte Dauber, Matthias Platzer, Sven M. Bergmann, Daniela Schrudde, Andreas Sauerbrei and Roland Zell

J. Virol. 2013, 87(19):10895. DOI: 10.1128/JVI.01094-13.

Published Ahead of Print 24 July 2013.

Updated information and services can be found at:
<http://jvi.asm.org/content/87/19/10895>

These include:

REFERENCES

This article cites 47 articles, 20 of which can be accessed free at: <http://jvi.asm.org/content/87/19/10895#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Characterization of a Novel Picornavirus Isolate from a Diseased European Eel (*Anguilla anguilla*)

Dieter Fichtner,^a Anja Philipps,^{b*} Marco Groth,^c Heike Schmidt-Posthaus,^d Harald Granzow,^a Malte Dauber,^e Matthias Platzer,^c Sven M. Bergmann,^a Daniela Schrudde,^a Andreas Sauerbrei,^b Roland Zell^b

Institute of Infectology, Friedrich Loeffler Institut, Federal Research Institute for Animal Health, Greifswald-Insel Riems, Germany^a; Department of Virology and Antiviral Therapy, Jena University Hospital, Friedrich Schiller University, Jena, Germany^b; Genome Analysis, Leibniz Institute for Age Research, Fritz Lipmann Institute, Jena, Germany^c; Centre for Fish and Wildlife Health, Institute of Animal Pathology, University of Bern, Bern, Switzerland^d; Institute for Virus Diagnostics, Friedrich Loeffler Institut, Federal Research Institute for Animal Health, Greifswald-Insel Riems, Germany^e

A novel picornavirus was isolated from specimens of a diseased European eel (*Anguilla anguilla*). This virus induced a cytopathic effect in eel embryonic kidney cells and high mortality in a controlled transmission study using elvers. Eel picornavirus has a genome of 7,496 nucleotides that encodes a polyprotein of 2,259 amino acids. It has a typical picornavirus genome layout, but its low similarity to known viral proteins suggests a novel species in the family *Picornaviridae*.

All members of the family *Picornaviridae* are small, nonenveloped, icosahedral viruses with positive-strand RNA genomes. Genome sizes range from 7,000 to 9,100 nucleotides. The 5' end of the RNA is covalently linked to a small virus-encoded peptide (3B); the 3' end is polyadenylated. The genome encodes one or two polyproteins that are processed co- and posttranslationally into 10 to 14 mature polypeptides. Cap-independent translation initiation is stimulated by one or two internal ribosome entry site (IRES) elements. Common to all picornaviruses are homologous capsid proteins and the nonstructural proteins 2C, 3C^{pro}, and 3D^{pol}. The sequences of other nonstructural proteins (leader protein, 2A, 2B, 3A, and 3B) are not conserved and may be unique to some picornavirus genera (1, 2).

The family *Picornaviridae* is currently comprised of 37 species grouped into 17 genera (*Aphthovirus*, *Aquamavirus*, *Avihepatovirus*, *Cardiovirus*, *Cosavirus*, *Dicipivirus*, *Enterovirus*, *Erbovirus*, *Hepatovirus*, *Kobuvirus*, *Megrivirus*, *Parechovirus*, *Salivirus*, *Sapelovirus*, *Senecavirus*, *Teschovirus*, and *Tremovirus*; <http://ictvonline.org/>). However, picornavirus taxonomy is still in flux, as next-generation sequencing technologies facilitate metagenomics studies using viral nucleic acid-containing specimens. Analysis of viral genomes without prior virus isolation led to the identification of more than 30 approved or tentative picornavirus species from humans (cosavirus [3], salivirus [4], scaffold virus [5], and rhinovirus-C [6]), rodents (mosavirus, rosavirus, and murine kobuvirus [7]), pinnipeds (seal picornavirus [8] and California sea lion sapelovirus [9]), domestic cats and dogs (feline picornavirus [10], canine picornavirus [11], canine kobuvirus [12], and canine picodicrovirus [2]), ungulates (bovine hungaravirus, ovine hungaravirus [13], swine pasivirus [14], and porcine kobuvirus [15]), birds (turdiviruses 1 to 3 [16], turkey hepatitis virus [17], pigeon picornaviruses A and B [18], quail picornavirus [19], and gallivirus [20]), and bats (bat picornaviruses 1 to 3 [21], bat kobuvirus-like virus [22], *Rhinolophus affinis* picornavirus, *Ia io* picornavirus, and *Miniopterus schreibersii* picornavirus [23]).

Although all published picornaviruses were identified in mammals and birds, there are reports of picornavirus-like viruses of reptiles (24, 25), fish (26–31), and marine invertebrates (32–35), indicating circulation of such viruses in aquatic ecosystems. Here, we describe a fish picornavirus which proves that picornavirus

ecology indeed includes lower vertebrates of freshwater and probably marine ecosystems. Eel picornavirus 1 (EPV-1) was isolated from a European eel (*Anguilla anguilla*) from Lake Constance on the Rhine River. The viral genome contains an open reading frame (ORF) of 6,777 nucleotides (2,259 amino acids [aa]), and the predicted polyprotein displays the typical organization of a picornavirus. According to the Picornavirus Study Group criteria (www.picornastudygroup.com/definitions/genus_definition.htm), low similarity to the conserved P1 and 3CD precursor proteins of EPV-1 suggests a novel genus in the family *Picornaviridae*.

Starting in May 2005, an increased number of diseased or dead eels were observed in fish traps and aquacultures in the Lake Constance area. Diseased fish presented with increased mucus production, mucous ulcers, and reddened skin but no hemorrhages. One sample (F15/05) that was sent to the National Reference Laboratory for Fish Diseases, Insel Riems, Germany, included specimens of heart, spleen, kidney, liver, and brain. In routine bacteriology by a regional laboratory, only *Aeromonas hydrophila* was detected. By PCR, the genome of the anguillid herpesvirus (*Herpesvirus anguillae* [HVA]) was detected in the kidney. A cytopathogenic, chloroform-resistant virus could be isolated from all four organ samples using eel embryonic kidney (EK-1) cells [catalogue no. CCLV-RIE 809, Collection of Cell Lines in Veterinary Medicine (CCLV) of the Friedrich Loeffler Institut]. This virus exhibited a picornavirus-like appearance in electron microscopy (Fig. 1A) and was designated EPV-1 F15/05. After passaging of the virus, HVA-specific PCR was negative. EPV-1 F15/05 was propagated in EK-1 cells. Infected cell cultures were incubated at 26°C. Three to 4 days postinfection (p.i.), a cytopathic effect was visible (Fig. 1B).

Received 22 April 2013 Accepted 3 July 2013

Published ahead of print 24 July 2013

Address correspondence to Roland Zell, Roland.Zell@med.uni-jena.de.

* Present address: Anja Philipps, Geneart—Life Technologies, Regensburg, Germany.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JVI.01094-13




FIG 1 Characterization of eel picornavirus. (A) Electron micrograph of EPV-1 from supernatant of infected EK-1 cells. Bar, 100 nm. (B) Cytopathic effect of EK-1 cells induced by EPV-1 after 3 days p.i. The inset shows uninfected cells. (C) Survival rates of infected and control eels. Glass eels ($n = 16$) were infected with EPV-1 by bathing for 1 h in virus-containing water. A control group of 12 glass eels was mock infected. The eels were kept in aquaria for 21 days. One glass eel of the control group died after an injury. (D) Histopathological changes in the liver of an experimentally infected glass eel. Hepatocytes are irregularly vacuolated (closed arrowheads). Multiple cell membranes are disrupted (open arrowheads). There are karyopyknosis (arrows)

As EPV-1 was isolated from an eel with a mixed infection, its virulence was examined in a controlled infection experiment. A group of glass eels (length, approximately 10 cm; $n = 16$) was infected by bathing in virus-containing water (10^6 50% tissue culture infective doses [TCID₅₀]/ml water) for 1 h. Subsequently, the eels were kept in 400-liter aquaria at a water temperature of 20°C for 21 days. A control group of eels ($n = 12$) was treated and kept under the same conditions and the same volume of an analogous cell culture medium without adding virus to the bath. Beginning 3 days p.i., infected eels died without external signs; at the end of the experiment, 7 of the 16 infected eels had died (43%) (Fig. 1C). In the untreated control group, one eel died in consequence of an injury. Histological examination of all organs, performed at the Centre for Fish and Wildlife Health, Bern, Switzerland, showed decreased numbers of fat vacuoles in hepatocytes of infected euthanized and dead eels compared to uninfected controls, increased single-cell necrosis, scattered small necrotic foci, and mild infiltrations of lymphocytes and macrophages in the liver parenchyma (Fig. 1D). In the kidneys, vacuolization and hyaline droplet degeneration of renal tubular epithelium and multiple small necroses in the interstitium were observed. Virus could be reisolated from 10 of 10 infected euthanized or dead eels and was identified as picornavirus by means of indirect immunofluorescence assay (Fig. 1E) and electron microscopy. No virus could be isolated from the control group.

The genome of EPV-1 was sequenced using the Illumina/Solexa method. Viral RNA was purified from EPV-infected EK-1 cells. Virus was released from infected cells showing cytopathic effect by three freeze-thaw steps. After pelleting of cell detritus (centrifugation at $4,000 \times g$ for 20 min), virus was sedimented by ultracentrifugation ($100,000 \times g$, 3 h, 4°C). RNA extraction from the pellets and sample preparation for Illumina sequencing were done as described previously (36). Sequencing was performed on a GAIIX (Illumina) instrument to create reads with lengths of 76 nucleotides (nt). For assembly, the reads were mapped to the NCBI nucleotide sequence database to identify similar picornavirus reads. These were used for *de novo* assembly using ABySS (37) with a k -mer length of 60. Four internal gaps of 73, 67, 60, and 34 nucleotides as well as the 3' end (approximately 180 nt) and the very 5' end were not covered by this approach (Fig. 2). The gaps and the 3' end were sequenced by conventional Sanger sequencing as described previously (36). All attempts to amplify the very 5' end by rapid amplification of 5' cDNA ends (5' RACE) failed.

As depicted in Fig. 2, the genome of EPV-1 exhibits a typical picornaviral genome organization and is comprised of 7,496 nucleotides with a 5' untranslated region (UTR) of at least 488 nt, a large ORF of 6,777 nt, and a long 3' UTR of 231 nt. The genome of EPV-1 has no obvious leader peptide-encoding gene region. Conserved Rhv domains (ranging from aa 130 to 245 and 485 to 630) were found in the capsid proteins 1AB and 1C (<http://pfam.sanger.ac.uk/>). The P1 precursor is most likely released by an aphthovirus-like 2A1 peptide. The 2A2 gene region exhibits the H-box/NC sequence motifs (H₇₉₇YGV and N₈₆₅CE), but the ALTXXKAXXX

and small areas of necrosis and fibrin exudation (stars). Bar = 25 μ m. Hematoxylin-eosin stain. (E) Detection of EPV-1 by indirect immunofluorescence using the rabbit antiserum T51 (in-house production) and goat anti-rabbit IgG-FITC fluorescein isothiocyanate (Sigma), with counterstaining of nuclei with propidium iodide. The inset shows uninfected cells.




FIG 2 Schematic depiction of the eel picornavirus genome. The 5' and 3' UTRs and the ORF (box) are displayed. Filled triangles show the 3C^{Pro} processing sites; the ribosomal skip sequence is indicated by a pound sign. (A)_n poly(A) tail. Nucleotide (nt) and amino acid (aa) positions of putative cleavage sites are given as well as the amino acid positions of the P1/P1' sites.

KXXL motif, which is common to the 2A proteins of parechoviruses, is modified (A₈₁₈TTXKVXXXXXXD). The 2B protein has homology to the 2B protein of parechoviruses (38), duck hepatitis A virus (DHAV-1) (39), and swine pasivirus (14) (Table 1). The 2C gene region exhibits three conserved sequence motifs (A, G₁₁₈₉XXGXGKS; B, Q₁₂₄₄YFHVIDDLAQ; C, K₁₂₈₃GMSYTSRVVIA TTN) (reference 40 and literature cited therein). 3A shares some similarity with Ljungan virus (LV) (41) but no obvious homology to any other known picornavirus (Table 1). The N terminus of 3B (RAY₁₅₇₀) is homologous to that of the parechovirus 3B. The putative 3C protease region encodes a GXC₁₇₄₉G active site and a G₁₇₆₅XH substrate-binding pocket (40). Conserved motives of the 3D polymerase are the K₁₉₅₄DELRL, G₂₀₈₁GMPSG, Y₂₁₂₂GDD, and F₂₁₇₀LKR sequences (40). Table 1 compares the predicted processing products of EPV-1 with those of its closest relatives. In general, EPV-1 exhibits low similarity to all known picornaviruses.

For phylogenetic analyses, the P1 gene region and the 3CD-encoding gene region of EPV-1 were compared to the corresponding gene regions of 64 picornaviruses representing all approved or tentative picornavirus species. Reference picornavirus sequences were retrieved from GenBank and used for comparisons and phylogenetic analyses. Nucleotide sequences were aligned with MEGA5 (42) and adjusted manually. For phylogenetic tree reconstruction, four Bayesian Metropolis-coupled Markov chains were calculated with MrBayes 3.1.2 (43) using an optimal substitution model. Modeltest implemented in MEGA5 suggested GTR+G+I

for both analyses. Convergence was reached after 674,000 generations (P1 region) and 322,000 generations (3CD region). Both phylogenetic trees revealed that *Parechovirus*, *Avihepatovirus*, *Aquamavirus*, and the swine pasivirus are the closest relatives of EPV-1. In addition, the P1 gene region of *Rhinolophus affinis* picornavirus is also related to EPV-1 (Fig. 3). Low amino acid identity as shown in Table 1 and Fig. 3 suggests that EPV-1 constitutes a novel picornavirus species and probably a novel picornavirus genus. Therefore, we propose the species name “*Anguavirus*.”

So far, few aspects of picornavirus ecology have been elucidated. Due to the fecal-oral transmission route, picornaviruses can be considered environmental viruses, and detection of enteroviruses, hepatitis A virus, and Aichi virus in shellfish (44, 45) as well as the identification of seal picornavirus (8) and an enterovirus from the bottlenose dolphin (46) confirms this view. However, PCR detection of picornaviruses does not necessarily indicate their replication in bivalve molluscs but suggests contamination with sewage and bioaccumulation. Viruses with picornavirus-like morphology in electron micrographs have been observed repeatedly in fish, reptiles, and invertebrates, but virus isolation and propagation in cultured cells have often failed (for example, see references, 24, 25, 28, 31, 32, 33, 35, 47, and 48). Therefore, isolation of EPV-1 from eels, experimental infection with induction of clinical signs, and reisolation constitute proof that picornaviruses indeed infect lower vertebrates and raise the question of how sta-

TABLE 1 Pairwise amino acid identities^a

EPV-1 protein ^b	% Identity with protein from:					
	<i>Parechovirus</i> HPeV-1 (L02971)	<i>Parechovirus</i> LV-1 (AF327920)	<i>Avihepatovirus</i> DHAV-1 (DQ249299)	<i>Aquamavirus</i> SePV-1 (EU142040)	Unassigned SPaV-1 (JQ316470)	Unassigned <i>R. affinis</i> PV-1 (JQ814853) ^c
1AB	30.1	33.6	25.3	27.7	24.5	17.1
1C	26.6	31.2	22.7	31.1	30.8	29.2
1D	22.8	22.7	18.7	15.9	21.5	21.5
2A1 ^d		24.0	52.2	20.7	36.4	40.0
2A2 ^e	27.9	33.8	30.0			
2B	28.6	25.8	21.3	13.3	19.7	
2C	27.6	27.3	29.9	26.5	27.3	
3A	8.8	20.3	12.9	11.0	12.2	
3C	18.3	19.1	21.0	26.1	19.1	
3D	34.6	41.6	38.6	29.0	31.3	

^a EPV, eel picornavirus; HPeV, human parechovirus; LV, Ljungan virus; DHAV, duck hepatitis A virus; SePV, seal picornavirus; SPaV, swine pasivirus; *R. affinis* PV, *Rhinolophus affinis* picornavirus.

^b Predicted proteins. 3B was excluded.

^c Partial sequence comprising 1AB, 1C, 1D, and 2A1.

^d Aphthovirus-like 2A peptide with cleavage at the NPG/P sequence.

^e 2A protein with H-box/NC motif.




FIG 3 Phylogenetic analyses of the P1-encoding (1AB-1D) (A) and the 3CD-encoding (B) gene regions of 65 picornavirus strains. Sixty-four sequences obtained from the GenBank and the EPV-1 sequence were included. The tree was inferred with MrBayes 3.1.2 using the GTR substitution model assuming gamma distribution and invariant sites. GenBank accession numbers, (tentative) species names, and strain designations are presented. The scale bar indicates substitutions per site. Numbers at nodes indicate posterior probabilities.

ble infection chains are maintained in aquatic ecosystems with high dilution and/or continuous outflow.

Nucleotide sequence accession number. The RNA sequence of EPV-1 F15/05 was submitted to GenBank with accession no. [KC843627](https://doi.org/10.1093/nuclemta/kty001).

ACKNOWLEDGMENTS

We thank Martina Müller and Ivonne Görlich for excellent technical assistance.

REFERENCES

- Knowles NJ, Hovi T, Hyypia T, King AMQ, Lindberg AM, Pallansch MA, Simmonds P, Skern T, Stanway G, Yamashita T, Zell R. 2012. Family Picornaviridae, p 855–880. *In* King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (ed), *Virus taxonomy*. Ninth report of the International Committee on Taxonomy of Viruses. Elsevier Academic Press, Amsterdam, The Netherlands.
- Woo PCY, Lau SKP, Choi GKY, Huang Y, Teng JLL, Tsoi HW, Tse H, Yeung ML, Chan KH, Jin DY, Yuen KY. 2012. Natural occurrence and characterization of two internal ribosome entry site elements in a novel

virus, canine picodicitrovirus, in the picornavirus-like superfamily. *J. Virol.* 86:2797–2808.

- Kapoor A, Victoria J, Simmonds P, Slikas E, Chieochansin T, Naem A, Shaikat S, Sharif S, Alam MM, Angez M, Wang C, Shafter RW, Zaidi S, Delwart E. 2008. A highly prevalent and genetically diversified Picornaviridae genus in South Asian children. *Proc. Natl. Acad. Sci. U. S. A.* 105:20482–20487.
- Li L, Victoria J, Kapoor A, Blinkova O, Wang C, Babrzadeh F, Mason CJ, Pandey P, Triki H, Bahri O, Oderinde BS, Baba MM, Bukbuk DN, Besser JM, Bartkus JM, Delwart EL. 2009. A novel picornavirus associated with gastroenteritis. *J. Virol.* 83:12002–12006.
- Jones MS, Lukashov VV, Ganac RD, Schnurr DP. 2007. Discovery of a novel human picornavirus in a stool sample from a pediatric patient presenting with fever of unknown origin. *J. Clin. Microbiol.* 45:2144–2150.
- Lau SK, Yip CC, Tsoi HW, Lee RA, So LY, Lau YL, Chan KH, Woo PC, Yuen KY. 2007. Clinical features and complete genome characterization of a distinct human rhinovirus (HRV) genetic cluster, probably representing a previously undetected HRV species, HRV-C, associated with acute respiratory illness in children. *J. Clin. Microbiol.* 45:3655–3664.
- Phan TG, Kapusinszky B, Wang C, Rose RK, Lipton HL, Delwart EL. 2011. The fecal flora of wild rodents. *PLoS Pathog.* 7:e1002218. doi:10.1371/journal.ppat.1002218.

8. Kapoor A, Victoria J, Simmonds P, Wang C, Shafer RW, Nims R, Nielsen O, Delwart E. 2008. A highly divergent picornavirus in a marine mammal. *J. Virol.* 82:311–320.
9. Li L, Shan T, Wang C, Cote C, Kolman J, Onions D, Gulland FMD, Delwart E. 2011. The fecal flora of California Sea lions. *J. Virol.* 85:9909–9917.
10. Lau SKP, Woo PCY, Yip CCY, Choi GKY, Wu Y, Bai R, Fan RYY, Lai KKY, Chan KH, Yuen KY. 2012. Identification of a novel feline picornavirus from the domestic cat. *J. Virol.* 86:395–405.
11. Woo PCY, Lau SKP, Choi GKY, Yip CCY, Huang Y, Tsoi HW, Yuen KY. 2012. Complete genome sequence of a novel picornavirus, canine picornavirus, discovered in dogs. *J. Virol.* 86:3402–3403.
12. Li L, Pesavento PA, Shan T, Leutenegger CM, Wang C, Delwart E. 2011. Viruses in diarrhoeic dogs include novel kobuviruses and sapoviruses. *J. Gen. Virol.* 92:2534–2541.
13. Reuter G, Pankovics P, Knowles NJ, Boros A. 2012. Two closely related novel picornaviruses in cattle and sheep in Hungary from 2008 to 2009, proposed as members of a new genus in the family *Picornaviridae*. *J. Virol.* 86:13295–13302.
14. Sauvage V, Gouilh MA, Cheval J, Muth E, Pariente K, Burguiere A, Caro V, Manuguerra JC, Eloit M. 2012. A member of the new *Picornaviridae* genus is shed in pig feces. *J. Virol.* 86:10036–10046.
15. Reuter G, Boldizsár A, Kiss I, Pankovics P. 2008. Candidate new species of kobuvirus in porcine hosts. *Emerg. Infect. Dis.* 14:1968–1970.
16. Woo PCY, Lau SKP, Huang Y, Lam CSF, Poon RWS, Tsoi HW, Lee P, Tse H, Chan ASL, Luk G, Chan KH, Yuen KY. 2010. Comparative analysis of six genome sequences of three novel picornaviruses, turdiviruses 1, 2 and 3, in dead wild birds, and proposal of two novel genera, *Orthoturdivirus* and *Paraturdivirus*, in the family *Picornaviridae*. *J. Gen. Virol.* 91:2433–2448.
17. Honkavuori KS, Shivaprasad LL, Briese T, Street C, Hirschberg DL, Hutchison SK, Lipkin WI. 2011. Novel picornavirus in turkey poults with hepatitis, California, USA. *Emerg. Infect. Dis.* 17:480–487.
18. Kofstad T, Jonassen CM. 2011. Screening of feral and wood pigeons for viruses harbouring a conserved mobile viral element: Characterization of novel astroviruses and picornaviruses. *PLoS One* 6:e25964. doi:10.1371/journal.pone.0025964.
19. Pankovics P, Boros A, Reuter G. 2012. Novel picornavirus in domesticated common quail (*Coturnix coturnix*) in Hungary. *Arch. Virol.* 157: 525–530.
20. Farkas T, Fey B, Hargitt E, Parcells M, Ladman B, Murgia M, Saif Y. 2012. Molecular detection of novel picornaviruses in chickens and turkeys. *Virus Genes* 44:262–272.
21. Lau SK, Woo PC, Lai KK, Huang Y, Yip CC, Shek CT, Lee P, Lam CS, Chan KH, Yuen KY. 2011. Complete genome analysis of three novel picornaviruses from diverse bat species. *J. Virol.* 85:8819–8828.
22. Li L, Victoria JG, Wang C, Jones M, Fellers GM, Kunz TH, Delwart E. 2010. Bat guano virome: predominance of dietary viruses from insects and plants plus novel mammalian viruses. *J. Virol.* 85:6955–6965.
23. Wu Z, Ren X, Yang L, Hu Y, Yang J, He G, Zhang J, Dong J, Sun L, Du J, Liu L, Xue Y, Wang J, Yang F, Zhang S, Jin Q. 2012. Virome analysis for identification of novel mammalian viruses in bat species from Chinese provinces. *J. Virol.* 86:10999–11012.
24. Essbauer S, Ahne W. 2001. Viruses of lower vertebrates. *J. Vet. Med. B Infect. Dis. Vet. Public Health* 48:403–475.
25. Heldstab A, Bestetti G. 1984. Virus associated gastrointestinal diseases in snakes. *J. Zoo Animal Med.* 15:118–128.
26. Ahne W, Anders K, Halder M, Yoshimizu M. 1990. Isolation of picornavirus-like particles from the European smelt, *Osmerus eperlanus* (L.). *J. Fish Dis.* 13:167–168.
27. Bloch B, Gravningen K, Larsen JL. 1991. Encephalomyelitis among turbot associated with a picornavirus-like agent. *Dis. Aquat. Org.* 10: 65–70.
28. Hetrick FM, Hedrick RP. 1993. New viruses described in finfish from 1988–1992. *Annu. Rev. Fish Dis.* 3:187–207.
29. Iwanowicz LR, Goodwin AE, Heil N. 2000. A small RNA virus isolated from apparently healthy wild sandbar shiners, *Notropis scepticus* (Jordan & Gilbert). *J. Fish Dis.* 23:349–352.
30. Moore AR, Li MF, McMenemy M. 1988. Isolation of a picorna-like virus from smelt, *Osmerus mordax* (Mitchill). *J. Fish Dis.* 11:179–184.
31. Sano T. 1995. Viruses and viral diseases of salmonids. *Aquaculture* 132: 43–52.
32. Ahne W. 1994. Viral infections of aquatic animals with special reference to Asian aquaculture. *Annu. Rev. Fish Dis.* 4:375–388.
33. Johnson PT. 1978. Viral diseases of the blue crab, *Callinectes sapidus*. *Mar. Fish. Rev.* 40:13–15.
34. Kuris AM, Poinar GO, Jr, Hess R, Morris TJ. 1979. Virus particles in an internal parasite *Portunio conformis* (Crustacea: Isopoda), and its marine crab host, *Hemigrapsus oregonensis*. *J. Invertebrate Pathol.* 34:26–31.
35. Rasmussen LPD. 1986. Virus-associated granulocytomas in the marine mussel, *Mytilus edulis*, from three sites in Denmark. *J. Invertebrate Pathol.* 48:117–123.
36. Philipps A, Dauber M, Groth M, Schirrmeier H, Platzer M, Krumbholz A, Wutzler P, Zell R. 2012. Isolation and molecular characterization of a second serotype of the encephalomyocarditis virus. *Vet. Microbiol.* 161: 49–57.
37. Biro I, Jackman SD, Nielsen CB, Qian JQ, Varhol R, Stazyk G, Morin RD, Zhao Y, Hirst M, Schein JE, Horsman DE, Connors JM, Gascoyne RD, Marra MA, Jones SJ. 2009. De novo transcriptome assembly with ABySS. *Bioinformatics* 25:2872–2877.
38. Hyypää T, Horsnell C, Maaronen M, Khan M, Kalkinen N, Auvinen P, Kinnunen L, Stanway G. 1992. A distinct picornavirus group identified by sequence analysis. *Proc. Natl. Acad. Sci. U. S. A.* 89:8847–8851.
39. Tseng CH, Knowles NJ, Tsai HJ. 2007. Molecular analysis of duck hepatitis virus type 1 indicates that it should be assigned to a new genus. *Virus Res.* 123:190–203.
40. Koonin EV, Dolja VV. 1993. Evolution and taxonomy of positive-strand RNA viruses: implications of comparative analysis of amino acid sequences. *Crit. Rev. Biochem. Mol. Biol.* 28:375–430.
41. Johansson S, Niklasson B, Maizel J, Gorbalenya AE, Lindberg AM. 2002. Molecular analysis of three Ljungar virus isolates reveals a new, close-to-root lineage of the *Picornaviridae* with a cluster of two unrelated 2A proteins. *J. Virol.* 76:8920–8930.
42. Tamura K, Peterson D, Peterson M, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28:2731–2739.
43. Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574.
44. Di Pinto A, Conversano MC, Forte VT, La Salandra G, Montervino C, Tantillo GM. 2004. A comparison of RT-PCR based assays for the detection of HAV from shellfish. *New Microbiol.* 27:119–124.
45. Hansman GS, Oka T, Li TC, Nishio O, Noda M, Takeda N. 2008. Detection of human enteric viruses in Japanese clams. *J. Food Prot.* 71: 1689–1695.
46. Nollens HH, Rivera R, Palacios G, Wellehan JFX, Saliki JT, Caseltine SL, Smith CR, Jensen ED, Hui J, Lipkin WI, Yochem PK, Wells RS, St Leger J, Venn-Watson S. 2009. New recognition of enterovirus infections in bottlenose dolphins (*Tursiops truncatus*). *Vet. Microbiol.* 139:170–175.
47. Glazebrook JS, Heasman MP, de Beer SW. 1990. Picorna-like viral particles associated with mass mortalities in larval barramundi, *Lates calcarifer* Bloch. *J. Fish Dis.* 13:245–249.
48. Yoshikoshi K, Inoue K. 1990. Viral nervous necrosis in hatchery-reared larvae and juveniles of Japanese parrotfish, *Oplegnathus fasciatus* (Temminck & Schlegel). *J. Fish Dis.* 13:69–77.